

COMMUNITY SPORTS COMMITTEE

Report of the Advisory Committee for the Healthy Exercise for All Campaign — Physical Fitness Test for the Community

Purpose

This paper seeks to report to Members of the Community Sports Committee (CSC) on the study report on the Physical Fitness Test for the Community (Project) under the Healthy Exercise for All Campaign and the recommendations on future promotion plans.

Background

2. The General Administration of Sport of China (GASC) has carried out a national physical fitness test every five years since 2000. People of different organisations were invited to participate in the test, which comprised sample surveys conducted on the physique, physiological functions and physical fitness of the populations of various provinces (including regions and municipalities), in order to determine the overall physical fitness condition of the whole population and identify the change patterns. The first territory-wide physical fitness test in Hong Kong was completed during the period from 2005 to 2006 under the guidance of the Advisory Committee for the Physical Fitness Test for the Community under the CSC. To tie in with the China National Physical Fitness Surveillance Project (全民體質監測計劃) of the GASC and take forward the recommendation of conducting a territory-wide physical fitness test every five years, the CSC endorsed the proposal to conduct the second territory-wide physical fitness test at the meeting on 17 March 2010 with a view to collecting data on the physical fitness of Hong Kong people aged 3 to 69. Through launching the project, we hope that individual participants will have a general understanding of their own physical fitness condition, while the information collected can continue to contribute to the database on the physical fitness of Hong Kong people, help to determine the correlation between the physical activity pattern and the physical fitness of Hong Kong people, and facilitate the identification of the

priority areas for improvement so as to enhance the general physical fitness of the public.

3. In parallel, the CSC endorsed the setting up of the Advisory Committee (AC) for the Healthy Exercise for All Campaign — Physical Fitness Test for the Community to give advice and recommendations on the work of the Project including the study proposal, publicity and implementation strategy as well as the preparation of study report . The AC is composed of representatives from the CSC (to serve as the convenor and deputy convenor of the AC respectively), the Leisure and Cultural Services Department (LCSD), Education Bureau, Department of Health, the Hong Kong Association of Sports Medicine and Sports Science, and the Physical Fitness Association of Hong Kong, China.

4. To ensure the smooth implementation of this large-scale Project, the following three professional bodies were engaged by the LCSD through open tender in early 2011 and charged with the technical work of the Project:

- (a) The Chinese University of Hong Kong — to provide advisory services on the project design, monitoring and co-ordinating services for project implementation, and statistical services for data analysis and compilation of study report.
- (b) Physical Fitness Association of Hong Kong, China — to undertake the study of the physical fitness of students under the Project; to conduct physical fitness tests for secondary school students (aged 13 to 19) and kindergarten children (aged 3 to 6); to provide statistical services for data analysis and compilation of study report; to conduct physical fitness test for adults (aged 20 to 59) and the elderly (aged 60 to 69); and to carry out data collection and input.
- (c) MVA Hong Kong Limited — to provide statistical survey services for adults (aged 20 to 59) and the elderly (aged 60 to 69).

5. In the project launched in 2005, the method of convenience sampling was adopted. Through publicity and invitation across the territory, the previous project involved participants from different organisations and groups as well as interested members of the public joining the fitness test on the spot. The findings of the study only reflected the physical fitness of the participants. With the experience gained

from the first physical fitness test and the professional views collected from the academic field, the AC proposed that the present Project should adopt a random sampling method to ensure representativeness of the findings and to reflect more accurately the physical fitness of Hong Kong people.

6. On this basis, the Project adopted the random sampling method in a comprehensive manner. Target participants were Hong Kong people aged between 3 and 69, categorised into five age groups: infants (aged 3 to 6), children (aged 7 to 12), adolescents (aged 13 to 19), adults (aged 20 to 59), and the elderly (aged 60 to 69). In the age groups of infants (aged 3 to 6) and adolescents (aged 13 to 19), one kindergarten and one secondary school were selected respectively from each of the 18 districts across the territory by random sampling to participate in the Project. Samples for children (aged 7 to 12) were collected through the “Survey Study of Students’ Physical Fitness and Their Attitudes towards Physical Education 2010/11” under the assistance of the Education Bureau. The physical fitness test and questionnaire survey for the three groups were conducted at schools/kindergartens. For the collection of samples of adults (aged 20 to 59) and the elderly (aged 60 to 69), the questionnaire survey was conducted by inviting eligible family members of the households selected from the 18 districts to complete a questionnaire. Those who had completed the questionnaire were then invited to take the physical fitness test in the sports centre near their homes. Data collection for the Project was launched in April 2011 and completed in January 2012, with over 13 000 samples collected in the questionnaire survey. The number of successful samples, in which the participants had completed both the questionnaire survey and physical fitness test, reached 8 178.

7. The Chinese University of Hong Kong (CUHK) analysed the data and reported the preliminary findings of the Project to Members at the 6th AC meeting held on 23 May this year. On the recommendations put forward by members, CUHK collected and analysed further information in order to refine the study report. At the 7th meeting held shortly on 13 June 2012, the AC examined and endorsed the final summary report (at [Annex](#)). The recommendations on plans for promotion were also discussed at the meeting.

Summary of Findings

8. The Project is the first of its kind in Hong Kong in which the territory-wide physical fitness test was conducted by random sampling. As the data obtained is representative of the physical fitness of Hong Kong people aged between 3

and 69, the findings of the study are worthy of the attention of all sectors of the community. The major findings are summarised as follows:

(a) ***Physical fitness level***

(i) **Overweight/Obesity**

Data of the study showed that approximately 12.9% of boys and 14.1% of girls among the infants (aged 3 to 6) were classified as overweight or obese according to the criteria set by the International Obesity Task Force. The proportion of overweight or obese in the infant group was considered relatively high and worthy of concern. Similarly, the obesity rate of children (aged 7 to 12) was relatively high with 30.6% of boys and 22.8% of girls being classified as overweight or obese. The proportion in the adolescent group (aged 13 to 19) was slightly lower with 18.5% of boys and 9.4% of girls being overweight or obese. For adults (aged 20 to 59) and the elderly (aged 60 to 69), 28.1% of men and 25.9 % of women were facing the problem of central obesity according to the waist circumference criterion. The problem was more serious in two age groups, namely the middle-aged men (aged 40 to 59) and the elderly women (aged 60 to 69), in which the obesity rates reached as high as 40% according to the classification of Body Mass Index ^{Note 1} (BMI). It was also found that the percentage of women with obesity increased with age. In view of the above, the problem of obesity should not be overlooked.

(ii) **Underweight**

The study also found that 19.4% of infants (aged 3 to 6) and 8.8% of children (aged 7 to 12) were underweight. As for adolescents (aged 13 to 19), 15.5% of boys and 21.0% of girls were underweight. In the adult group (aged 20 to 59), 14.0% of young women (aged 20 to 39) were underweight. In general, the underweight rate is higher in women than in men.

(iii) **High blood pressure**

Note 1 The classification of Body Mass Index (BMI) for Asians is recommended by the World Health Organisation (WHO) as follows –

Underweight: BMI < 18.5; Normal weight : 18.5 ≤ BMI < 23;

Overweight: 23 ≤ BMI < 25; Obesity : BMI ≥ 25

$$BMI = \frac{\text{Weight(kg)}}{\text{Height(m)}^2}$$

Our data showed that there was an increase in prevalence of hypertensive response (i.e. with the Resting Systolic BP \geq 140 mmHg or Resting Diastolic BP \geq 90 mmHg) from young to old adulthood. As young as 20 to 39 years old, 16.4% of men and 6.3% of women were already found at risk of high blood pressure. With the progression of age, the rate of risk for older men was approximately 2.4 times that of young men, and the rate for women was as high as 5.6 times.

Besides, in the children group, 4.1% of primary school students were identified as at risk of hypertension. Although the prevalence is low, caution should be taken as such problems are not expected in children at this young age.

(iv) Effects of gender and age on physical fitness

In the age groups of adults (aged 20-59) and the elderly (aged 60-69), men generally performed better than women in exercise performances (e.g. hand grip test, 1-min sit-up test, step test), except for the sit-and-reach test. Fitness levels decreased with age significantly in both genders, with women demonstrating a quicker decline.

(v) Effect of smoking status on physical fitness

The physical fitness of male non-smokers was significantly better than that of male smokers. For example, male non-smokers possessed smaller waist-to-hip ratio, greater leg power and balancing-controlling ability, and shorter reaction time. In females, as the relatively small number of female smokers in the samples has led to a relatively higher error rate in sampling, we cannot ascertain whether there is any significant difference between the physical fitness of smokers and non-smokers in women.

(vi) Correlation between educational attainment and physical fitness

In general, women with higher levels of education had better physical fitness, including smaller BMI, percent body fat, chest circumference, waist circumference, hip circumference and waist-to-hip ratio; lower resting BP; greater muscular strength; better cardio-respiratory fitness and balance-controlling ability, as well as shorter reaction time. Their higher level of physical fitness may be attributed to their consciousness of health and fitness.

For men, the correlation between educational attainment and physical fitness was less significant than women. The relatively significant differences mostly involved the comparison between the groups that had attained secondary and tertiary education respectively. In short, men with tertiary level of education generally had better physical fitness than men with secondary level of education, which was reflected in smaller BMI and percent body fat, lower Resting Diastolic BP, as well as greater abdominal endurance and leg strength.

(b) ***Physical activity pattern***

(i) **Physical activity participation**

Our data showed that less than 20% (17.7%) of infants (aged 3-6) spent one hour or more a day on outdoor activities. Apart from this, while over 80% (81.1%) of children (aged 7-12) perceived that they had adequate or very adequate physical activities as expressed in the questionnaires, only 9.5% and 7.0% of boys and girls respectively were classified as physically active according to the classification of physical activity participation by age group in the “2008 Physical Activity Guidelines for Americans” (American Indicator)^{Note 2} generally adopted in the academic field. There is a large discrepancy between the two sets of figures. In the group of adolescents (aged 13-19), boys were found to be more active than girls with 12.8% of boys and only 4.2% of girls being classified as physically active. In the age groups of adults (aged 20-59) and the elderly (aged 60-69), only less than 30% (28.6%) of participants were categorised as having adequate physical activity participation. Moreover, for every two adults in Hong Kong, one was found sedentary. Our data analysis also indicated that sedentary people had a relatively lower physical fitness level.

On the other hand, in the “Consultancy Study on Sport for All — Participation Patterns of Hong Kong People in Physical Activities” (the 2009 Consultancy

Note 2 “2008 Physical Activity Guidelines for Americans”: “Active” children are defined as those who have an accumulation of at least 60 minutes of physical activities of moderate or above intensity per day for 7 days a week. “Active” adolescents are defined as those who have an accumulation of at least 60 minutes of physical activities of moderate or above intensity per day for 7 days a week, including physical activities of vigorous intensity 3 days a week. “Active” adults are defined as those who have an accumulation of at least 150 minutes of physical activities of moderate or above intensity accumulated in a week (or having at least 75 minutes of physical activities of vigorous intensity, or a combination of physical activities of different intensity, e.g. 90 and 30 minutes of physical activities of moderate and vigorous intensity respectively).

Study) completed in 2009, the LCSD suggested that a “Baseline Indicator”^{Note 3}, which was encouraging, easier to meet and applicable to all age groups, should be adopted as a starting point to promote participation in physical activities. The targets would be the “inactive” and “some active” people. They were recommended to engage in physical activities of moderate or vigorous intensity for an accumulation of at least 30 minutes a day, at least three days a week. If the “Baseline Indicator” is used to assess the level of physical activity in the groups of children, adolescents, and adults and the elderly in the Project, only 51.8% (excluding physical education lessons), 42.3% and 37.1% of people in the respective groups can meet the indicator. The results are also unsatisfactory, showing that the levels of physical activity of all age groups are found inadequate.

(ii) Comparison of the distribution of data on the physical activity levels between the Project and the 2009 Consultancy Study (according to the classifications based on the Baseline Indicator and the American Indicator)

The data of the Project was compared and analysed according to the classifications based on the Baseline Indicator used in the 2009 Consultancy Study and the American Indicator, with an aim to examine the changes in the physical activity level of Hong Kong people (except the infant group) over the past few years. The 2009 Consultancy Study has been used for comparison mainly for its similarity with the Project in study methodology. Both studies adopted a random sampling method with the participants being categorised into the same age groups. In addition, the physical activity levels were assessed in a similar way. The findings of the comparison are as follows:

- Classification based on the Baseline Indicator

Our data showed that the percentage of adolescents classified as “active” dropped from 48.6% to 42.1%, mainly due to the significant decline in the percentage of “active” boys, which decreased from 57.5% to 50.1%. For adults and the elderly, the percentage of “active” participants fell from 43.5% to 37.0%, in which the percentages of men and women classified as “active” declined by 8.7% and 4.2% respectively, indicating a significant drop in both sexes.

Note 3 “Baseline Indicator”: For all age groups, “active” means having an accumulation of 30 minutes of physical activities of moderate or above intensity per day for at least 3 days a week.

- Classification based on the American Indicator

The percentage of children classified as “active” increased from 5.7% to 9.0%, mainly due to a significant rise in the percentage of girls (from 3.7% to 7.7%) being classified as physically active. There was no significant change in the physical activity level of adults and the elderly over the past few years, during which the percentage of the participants classified as “active” lied in the range of 28% to 30%. For the adolescent group, no comparison could be made because of the substantial differences in the questionnaires being used in the two studies.

As shown in the above data, the overall physical activity level of Hong Kong people has not shown any significant increase but rather a slight decline (except for girls aged 7 to 12) over the past few years. It is possible that the decline is the result of an unfavourable comparison against the data collected by the 2009 Consultancy Study when the public had a relatively higher awareness of the importance of physical exercise as some major multi-sports games such as the Equestrian Events of the 2008 Olympic Games and the 2009 East Asian Games were staged in Hong Kong around that time.

(iii) **Most participated physical/sports activities**

“Playing in the playground” (80.4%) and “swimming” (45.3%) were the most popular activities among infants. For children, 80% of participants attended sports training as an extra-curricular activity, in which ball games were the most popular sports chosen by 68.1% of the participants, followed by swimming (48.8%) and track and field (42.0%). In the adolescent group, ball games (75.1%), ice skating/roller skating (24.8%) and swimming (24.7%) were the most popular sports. For adults (aged 20-59) and the elderly (aged 60-69), ball games (40.4%) and running/jogging (36.9%) were the most popular sports among young adults. With the increase of age, walking (51.4%) took over the lead.

(iv) **Main reasons for and barriers to participation in physical activities**

The top three reasons for children to participate in sports activities were “I like physical activity”, “physical activity is very interesting” and “physical activity participation makes me feel happy” (with 4 as the full mark, the above three

reasons scored 3.61, 3.53 and 3.52 respectively). This showed that physical activities had to be interesting and effective in lifting the mood and bringing joyful feelings to the participants in order to attract children's participation. As for adolescents, "raising ability on sports", "health strengthening/prevention or cure of sickness", "releasing pressure and emotion" and "keeping fit" were the primary reasons for participating in physical activities (like distance running and weight training).

The top three barriers for children to participate in physical activities were "bad weather", "too much homework" and "being too tired" (with 3 as the full mark, the above three barriers scored 2.01, 1.99 and 1.86 respectively) whereas those for adolescents were "not interested" (52.3%), "lazy" (45.2%) and "lack of spare time" (44.3%). The questionnaire survey also found that over thirty percent (35.8%) of students were occasionally or regularly advised by their parents to reduce their physical activity participation in order to save time for study. For adults and the elderly, the three main barriers to sports participation were "lack of spare time" (62.9%), "tired" (37.9%) and "lazy" (33.5%).

(v) Influence of parents on children's participation in physical activities

Only 48.5% of infants (aged 3 to 6) participated in physical activities with their families at least once a week at weekends or on holidays in the past year. Another 31.8% of infants participated in physical activities with their families once or twice a month. The study revealed that children of parents with higher physical activity levels were relatively more active. In the adolescent group, statistics showed that the physical activity level of fathers had significant influence on the activeness of both their sons and daughters whereas that of mothers had obvious influence on their daughters only. Nevertheless, more than half (58.1%) of the adolescents said their families did not participate in any physical activities together at weekends or on holidays in the past year.

(vi) Correlation between other factors and physical fitness and physical activity level

In the groups of adults and the elderly, the higher the income level of the households, the more active the men were. Likewise, there was a close correlation between the type of occupation and physical activity level. Among the male respondents, 47.9% of "professionals" were categorised as "active",

which was the highest in all types of occupations. The greatest number of active females was found in the “managers or administrators” category, with 34.8% being classified as “active”.

(c) ***Survey findings regarding the “Sport For All Day”***

To sustainably promote “Sport for All” in the community and tie in with the National Fitness Day, the LCSD has organised a “Sport For All Day” every August since 2009. A series of free recreational and sports programmes are held in various districts for public participation. Since 2010, a number of fee-charging leisure facilities have also been provided for free public use with an aim to encourage regular participation in sports and other physical activities and enhance public awareness of the benefits and importance of doing exercise. To gauge the levels of awareness and involvement of the “Sport For All Day” among the public, relevant questions (three in total) were included in the questionnaires for the age groups of adolescents (aged 13 to 19) and adults and the elderly (aged 20 to 69) in the Project. It was found that 45.2% of adolescents and 46.4% of adults and the elderly had heard about the “Sport For All Day” organised by the LCSD. Among them, 32.3% of adolescents and 12.2% of adults and the elderly said they had taken part in the free recreational and sports programmes or made use of the free leisure facilities on the “Sport For All Day”. Of these participants, 67.4% of adolescents and 60.4% of adults and the elderly agreed that they had become more interested in engaging in sports and physical activities after their participation in the “Sport For All Day”.

Recommendations

9. The recommendations made in the report, based on the above study findings, are summarised as follows:

(a) ***On increasing the physical activity level of infants***

With only less than 20% of infants engaged in outdoor activities for one hour or more a day on average, the physical activity level of this group was rather low. It is recommended that parents should spend at least one hour a day on outdoor parent-child activities as far as possible. Kindergartens may incorporate one hour of physical games into their daily teaching programmes. Alternative mixes of the above two types of activities can also serve the purpose.

Furthermore, training on helping infants to acquire the correct physical movements (skills) should be offered to teachers. Government departments or stakeholders in the community may consider organising more workshops and fun days to encourage parents to carry out more family-based physical activities and teach them the importance of the activities.

(b) ***On providing programmes and facilities that meet the needs***

Stakeholders should, by making reference to the physical activities/sports that are most participated or popular in different age groups as stated in this report, organise more recreational and sports programmes and provide more facilities (ball pitches, swimming pools, indoor game rooms, children's playgrounds and jogging trails, etc.) that can meet the needs of the people of different ages and allow them to take part in these activities in their neighbourhood.

(c) ***On enhancing the awareness of physical activity level***

The data revealed that the physical activity level of more than half of the school children was inadequate. They spent considerable time on sedentary activities, such as watching television programmes and browsing the Internet. It is therefore recommended that children should be taught the concept of adequate physical activity levels since they start school. Relevant concepts on the frequency, duration and strength of physical activity should be covered to help them form a habit of doing regular exercise at a young age and build a healthy lifestyle.

Moreover, only 37.1% of families participated in regular physical activities together at weekends or on holidays in the past year. In view of the key role of parents in their children's participation in sports and physical activities, they should be encouraged to make good use of their leisure time to participate in physical activities together with their children. Stakeholders in the community may also design and organise more creative and innovative parent-child activities with the aim of increasing children's participation in physical activities.

(d) ***On publicising the "Baseline Indicator" of physical activity level***

The findings showed that the physical activity level of the general public of

Hong Kong was still far from adequate. Therefore, it is recommended that publicity on the “Baseline Indicator” should be continued. The public should be encouraged to participate in physical activities of moderate or vigorous intensity for an accumulation of at least 30 minutes a day, at least three days a week. This indicator, relatively attainable for those whose physical activity level is inadequate, can help them develop a habit of doing regular exercise. People who have already met this target may gradually increase their physical activity level towards the “American Indicator”.

(e) ***On relieving pressure and promoting adolescents’ interest in physical activities***

The findings shed light on the increasing study pressure faced by adolescents, which is probably related to their pursuit of academic achievement. The pressure accounts for the decline in their level of physical activity. Some academic researches reveal that school children having adequate physical activities tend to perform better in learning. As such, it is recommended that promotion campaigns should be organised to publicise the benefits of physical activities in enhancing academic performance through improvement in mental fitness in order to motivate students to participate in more physical activities. Meanwhile, to remove the main barriers for adolescents to participating in sports (“Not interested”), it is recommended that a diversified range of activities, such as cycling, dinghy sailing, canoeing and bowling, which are currently popular among young people, should be promoted in a sustained manner. Moreover, school teachers should be encouraged to add game elements to physical education classes so that even some dull physical fitness training can be turned into lively and interesting activities that appeal to students

(f) ***On promoting health information on diet and exercise and conducting regular physical fitness evaluations***

In view that there is a steady upward trend in overweight/obesity among children and adults, and that such risk factors as hypertension are diagnosed at younger ages, health information on various topics, including exercise training principles and health risks posed by physical inactivity, should be disseminated to the general public and updated regularly through various publicity channels ranging from thematic exhibitions, websites to leaflets. For example, all age groups should be encouraged to learn how to calculate their BMIs and be aware of the

classifications of overweight/obesity and central obesity. This can help the public to have a better understanding of their own physical fitness condition and adjust their physical activity levels accordingly. The public should also be encouraged to measure their blood pressure regularly and participate in physical fitness tests conducted at schools or organised by government departments so as to have a comprehensive understanding of their physical fitness condition and take appropriate improvement measures. Moreover, continuous efforts should be made to promote and institutionalise the implementation of food nutrition labelling, the standardisation of recommendations on daily nutrition intake by professional bodies, and the recommendation on exercise training programmes by qualified physical fitness instructors.

(g) ***On providing more opportunities for women to participate in sports***

The findings indicated that the physical fitness of women in general deteriorated with age faster than that of men. There was even an upsurge in the percentage of women suffering from hypertension. Nowadays women have to face different challenges and a lot of stress as they are playing a vital role in different aspects of society, including family, workplace and community. They need to have a strong physique in order to cope well with the heavy workload and different types of stress. It is recommended that consistent efforts should be made to strengthen the promotion of sports participation and the importance of developing a healthy lifestyle for women. On the one hand, women should be provided with more opportunities to participate in sports, and on the other hand, stakeholders in the community should be encouraged to organise more recreational and sports programmes that are better received by women, such as dancing, hydro fitness exercise, Tai Chi, and fitness exercises using fit ball and elastic band. This can encourage women's active participation in sports and help them maintain a good physique. Moreover, simple physical activities/sports that can be carried out at home or workplace should also be designed for them.

(h) ***On organising and promoting suitable activities for the middle-aged***

The findings of the study revealed that the problem of obesity among the middle-aged (aged 40 to 59) was quite serious. As most of the middle-aged are busy at work, it will be difficult for them to spare time for sports activities that are subject to the constraints of venue, time and number of participants. To

address the above, it is recommended that stakeholders should organise and promote activities that only require simple equipment and can be performed without time constraint, such as jogging and fitness walking. Other activities like gateball, lawn bowling, hydro fitness exercise and relaxation techniques are also suitable for the middle-aged, who can relieve their stress at work and in their daily life through more active participation in physical activities during their leisure time. Furthermore, it is recommended that the business sector may consider designing a sports area for their staff in the workplace and employing qualified instructors to teach their staff how to do appropriate exercise to help them experience the benefits of exercise. It is hoped that they will form a habit of doing exercise actively and regularly in their work environment.

The Way Forward

10. The study provides abundant physical fitness data and important reference for government departments and local organisations in health promotion. We suggest that future promotion should focus on disseminating the information and recommendations contained in the study report to stakeholders in the community so as to provide reference for them in formulating targeted policy in future. In parallel, the findings of the study should be published so as to allow the general public to understand the current physical fitness level of Hong Kong people as well as the importance of regular participation in sports and physical activities. In sum, the recommended promotion plan should cover the following four areas:

- (a) The key points of the study report and the related recommendations should be publicised through various media such as a press conference and a newspaper supplement to enable the public to have an understanding of the physical fitness level of the people of Hong Kong.
- (b) The study report should be released to the relevant government departments like the Department of Health and Education Bureau and uploaded onto the website of the LCSD for browsing and download by all organisations, groups and members of the public. Stakeholders in the community should also be encouraged to make reference to the relevant information and recommendations when formulating targeted policy, such as providing appropriate recreational and sports facilities and right combination of recreational and sports programmes for the public, and providing the public with more opportunities for participation in physical activities in order to help them improve their

physical fitness. The major targets include the following:

- District Councils (DCs)

The DCs perform the essential function of promoting community sports development. Serving as a bridge in the community, they provide key local ties to facilitate the co-ordination between district sports associations and community sports clubs in their respective districts. The DCs may, with reference to the study report and its recommendations, plan targeted recreational and sports programmes and facilities, and make continuous efforts to promote suitable recreational and sports programmes for different age groups, different sectors of the community and people with different abilities in order to encourage every member of the community to participate in physical activities and develop a strong physique.

- Schools

The findings of the study revealed that school children spent considerable time on sedentary activities (such as watching TV and browsing the Internet) and only 37.1% of the respondents had regular physical activities with their families at weekends or on holidays in the previous year. The data also showed that less than 20% of infants spent one hour or more a day on outdoor activities on average, reflecting a very inadequate level of physical activity among school children. Parents play an indispensable role in their children's participation in physical activities. As schools serve as an ideal platform for communication with parents, it is suggested that schools should put more emphasis on physical and sports activities and promote to parents the positive effects of physical activities on learning ability and academic performance of school children.

On the problem of inadequate physical activity among school children, pre-primary institutions may make reference to the Guide to the Pre-primary Curriculum (2006) and Reference Materials on Physical Activities in Kindergartens (1999) published by the Education Bureau as well as the Physical Activity Guide for Children Aged 2 to 6 (2011) published by the Department of Health so as to fully utilise the school hours to provide young children with more opportunities to participate in physical activities. At primary and secondary levels, schools may make reference to the Physical

Education Key Learning Area Curriculum Guide (2002) and An Introductory Guide to Fundamental Movement (2007) published by the Education Bureau to develop a comprehensive physical education curriculum covering a variety of activities in different areas. Through provision of diversified learning experiences, schools can help broaden students' choices and nurture students to participate regularly in at least one life-long physical activity, thereby helping them to develop a positive, active and healthy lifestyle.

● Government Departments and Public and Business Sectors

Government departments and public and business sectors may make reference to the findings of the study to devise a long-term promotion plan for recreation and sports development, education, health promotion and so on. Sustained efforts may also be made to organise and promote diversified recreational and sports programmes such as parent-child activities for all family members, recreational and sports activities popular among young people or women, and simple physical/sports activities that can be carried out at workplace. More recreational and sports facilities related to the above activities should also be provided to meet the demand of different groups.

Moreover, it is suggested that various organisations should call on their staff to do more exercise and encourage people having a low level of physical activity to adopt the “Baseline Indicator” as their starting point to participate in physical activities, i.e. participating in physical activities of moderate or vigorous intensity for an accumulation of at least 30 minutes a day, at least three days a week. Those who have already met the level may raise their physical activity level progressively and work towards the goal of achieving the “American Indicator” that suggests a higher level of physical activity, thereby developing a good habit of regular exercise.

- (c) Various publicity channels, including publishing brochures targeting different reader groups, should be used to raise public awareness of the physical fitness condition of their respective age groups. Designing a dedicated webpage on the LCSD website, producing a CD-ROM on the “Healthy Exercise for All Campaign”, organising roving exhibitions and arranging physical fitness test days should also be included in order to provide the general public with the guidelines on appropriate levels of exercise, indicators on participation in physical activities and relevant nutrition information. Through these efforts, the public can be

encouraged to exercise regularly in order to maintain a good physique and develop a healthy lifestyle.

- (d) Data on the physical fitness condition of the public should be collected on a regular basis to facilitate the continuous evaluation of the changes in their fitness condition. It is suggested that a territory-wide physical fitness test should be conducted every five years.

Advice Sought

- 11. Members are invited to comment on the above findings, recommendations and promotion plan.

Secretariat of the Advisory Committee for the
Healthy Exercise for All Campaign — Physical Fitness Test for the Community
July 2012

Healthy Exercise for All Campaign – Physical Fitness Test for the Community

Final Summary Report

Commissioned by

Community Sports Committee
of the Sports Commission

Co-ordinated by



Leisure and Cultural
Services Department

Supported by

Education
Bureau



Department of
Health



Hong Kong
Association of
Sports Medicine
and Sports
Science



Physical Fitness
Association of
Hong Kong,
China

Report submitted by



The Chinese University of Hong Kong
Department of Sports Science & Physical Education

【 Summary of Findings 】

1. Background

- 1.1 The Physical Fitness Test for the Community (the Study) was one of the events under the Healthy Exercise for All Campaign. The Study was steered by the Community Sports Committee of Hong Kong, and organized by the Leisure and Cultural Services Department (LCSD). It included a series of standardized physical fitness tests and questionnaire survey on the basis of “國民體質監測工作方案” done by the General Administration of Sport, China (國家體育總局).
- 1.2 Targeted Participants were Hong Kong citizens aged 3–69. There were five age groups, infants aged 3–6, children aged 7–12, adolescents aged 13–19, adults aged 20–59, and elderly aged 60–69.
- 1.3 The Study sampled over 13,000 participants through Stratified and random sampling method in 18 districts, ended up with 8,178 valid samples.
- 1.4 The LCSD as the organizer of the Study appointed the Department of Sports Science and Physical Education (SSPE) at The Chinese University of Hong Kong, the Physical Fitness Association of Hong Kong, China (HKFPA), and MVA Hong Kong Limited (MVA) to conduct a territory-wide community fitness survey from April 2011 to January 2012. The SSPE research team was to assist the Government Representative in monitoring the performance and provide consultation and professional advice on the work of other involved parties in the Project, and oversee the sampling method. Recommendation was made on fitness test and survey questionnaire design, including suggestions and modifications on testing protocols and questionnaire items, together with the format of data collection process. The SSPE was also responsible for data processing, statistical analysis, and the results report write-up. The HKFPA was responsible for testers training, conducting the fitness tests for all age groups, preparing and collecting questionnaire data from school-age groups, providing operational supporting services including safety screening and professional advice provision. MVA was responsible for enumerators training, conducting household interviews and inviting eligible citizens to the sports centers nearby for physical fitness tests.

1.5 There was a phased data collection process according to age groups. For infants and adolescents, the working group randomly selected one kindergarten and one secondary school in 18 districts respectively. For children, 18 primary schools were randomly selected from four regions (H.K. Island, Kowloon, New Territories East, and New Territories West) according to their modus operandi (direct subsidy scheme, aided school, and private independent school) with assistance from the Education Bureau (EDB). HKPFA was responsible for the data collection of the three student groups. For adults and elderly, MVA and HKPFA were to conduct household survey and follow up the fitness measurements respectively.

2. Project Objectives

2.1 The objectives of the Study were to (1) enable the individuals participating in the Physical Fitness Test to have a general understanding of their own physical fitness condition; (2) continue building up a database of the physical fitness of Hong Kong people; (3) recognize the relationship between the physical exercise pattern and the physical fitness of Hong Kong people; and (4) identify priority areas for improvement so as to enhance the overall physical fitness of the public.

2.2 Fitness measurements were executed by qualified HKPFA testers. The fitness data was used to generate normative tables by gender and age group, so that people could compare their fitness performance with the overall profile on their own.

2.3 The information about the patterns and the level of physical activity participation, lifestyle, demographic information as well as attitude towards participation in physical activity and self-perception of health-related issues were collected via questionnaire.

2.4 According to the findings, suggestions concerning education and related supporting policies were provided so as to raise Hong Kong people's physical activity level and general fitness profile.

3. Data Analysis

3.1 After all data were collected from questionnaire surveys and physical fitness tests, the research team carried out statistical analyses, including descriptive statistics, sampling distribution, comparisons, and correlation analysis. The relationships between participants' fitness levels and their exercise patterns as well as lifestyle habits and so on were investigated.

- 3.2** In the Study report, participants were being categorized into four groups (obese, overweight, underweight, and normal) according to their body mass index (BMI). The classification of adults and elderly was based on the BMI criteria for Asian established by the World Health Organization (WHO) in 2000 while infants, children, and adolescents were classified with reference to age and gender specific BMI criteria recommendation by the International Obesity Task Force (IOTF) in 2000 and 2007.
- 3.3** On the other hand, the Study grouped the participants aged 7 years or above according to their physical activity level. Two reference criteria were adopted, namely “2008 Physical Activity Guidelines for Americans” (“American Indicator”)¹ and “Physical Activity Baseline Indicator” (“Baseline Indicator”)² which was promoted locally and by some other countries/ regions, including China, England, and Taiwan. Participants were evaluated whether or not they met the recommendations under the two criteria. Although there was a difference in physical activity recommendations, research team still believed that it was necessary to conduct the classification analysis based on both criteria. The reasons included (1) “American Indicator” which was widely admitted and used in the field of physical education, medicine, and public health was founded on an enormous number of scientific evidences. It was quoted by WHO (2010) as well. The classification using a globally-recognized reference made helped the report more comparable; (2) “Baseline Indicator” which was less demanding than “American Indicator” and easily accepted by busy Hong Kong people was promoted locally in recent years. Therefore, research team summarized that each reference criteria had its advantages, especially in a fast-moving place like Hong Kong. People were encouraged to take a first step to meet “Baseline Indicator”. After enriching cognitive level as well as physical fitness level, they were suggested to take reference from “American Indicator” and gradually increase their physical activity level.

Remarks 1 “American Indicator”: “Active” children were those who accumulated at least 60 minutes moderate-or-above intensity physical activity every day in a week. “Active” adolescents were those who accumulated at least 60 minutes moderate-or-above intensity physical activity every day in a week in which any three days involved with vigorous intensity physical activity. “Active” adults and elderly were those who accumulated at least 150 minutes moderate-or-above intensity physical activity in a week (or 75minutes vigorous intensity physical activity or other combinations of different intensities, e.g. accumulation of 90 minutes and 30 minutes moderate- and vigorous- intensity physical activity respectively).

Remarks 2 “Baseline Indicator”: The Indicator was applied to all age groups. “Active” people were those who participated in moderate-or-above intensity physical activity at least three days a week with accumulation of 30 minutes or above per day.

- 3.4** The physical activity level of Hong Kong people was also used to compare with the one released in the “Report for the Consultancy Study on Sports for All – Participation Patterns of Hong Kong People in Physical Activities” in 2009 (“Sports for All Study 2009”). The purpose of making such comparison was to understand the change of physical activity level across the past few years. The main reason why “Sports for All Study 2009” was chosen for comparison was that both studies adopted similar research methodology: (1) random sampling; (2) same age groups of participants; (3) similar assessment methods of physical activity level.
- 3.5** The percentages of different parameters presented in the report were calculated from the true value before rounding. Therefore, add-ups of figures in the tables may not be equal to the sub-total or total.

4. Infants

4.1 Participants

- 4.1.1 One kindergarten was selected from 18 districts respectively through stratified randomly sampling. Each school assisted in picking one class from three grades respectively. Infants aged 3–6 were invited to participate in the Study.
- 4.1.2 There were 707 randomly selected samples of which 584 were valid (82.6%).
- 4.1.3 The research team compared the sample sizes of four sub-groups (boys/ girls aged 3–4; boys/ girls aged 5–6) with the “Mid-2011 Provisional Population Figures” published by the C&SD so as to allow necessary weighting calculations.

4.2 Physical Fitness Level

4.2.1 The mean values of body composition and fitness parameters were shown in the table below.

Age	3–4		5–6	
Parameters	Boys	Girls	Boys	Girls
Height (<i>cm</i>)	104.37	103.30	114.55	113.27
Weight (<i>kg</i>)	17.28	16.40	21.00	20.22
BMI (<i>kg/m²</i>)	15.83	15.33	15.91	15.66
Seated height (<i>cm</i>)	58.92	58.15	63.47	62.79
Total Skinfold (Upper Arm + Calf) (<i>mm</i>)	16.01	17.16	16.88	18.92
Chest circumference (<i>cm</i>)	54.73	53.94	58.30	57.35
Resting HR (<i>bpm</i>)	99.14	97.52	93.91	94.85
Sit-and-reach (<i>cm</i>)	5.35	7.52	4.42	6.39
Walking on a balance beam (<i>s</i>)	17.58	18.18	8.02	8.52
Continuous jump with both feet (<i>s</i>)	11.42	11.35	7.58	7.69
Standing long jump (<i>cm</i>)	67.00	62.93	90.98	81.89
10m × 2 shuttle run (<i>s</i>)	8.15	8.46	6.88	7.04
Throwing a Tennis Ball (<i>m</i>)	3.42	2.80	5.01	4.38

4.2.2 According to the criteria suggested by International Obesity Task Force (IOTF), 12.9% of boys and 14.1% of girls were classified as overweight or obese (see the table below). In addition, about 20% of infants (boys: 19.3%; girls: 19.5%) were categorized as underweight. To sum up, the prevalence of overweight and obese infants was a bit high. Close attention should be paid to the issue.

Age	Underweight		Overweight		Obese	
	Boys (%)	Girls (%)	Boys (%)	Girls (%)	Boys (%)	Girls (%)
3–4	20.1	22.2	6.7	9.0	3.4	0.7
5–6	18.4	16.5	8.2	13.5	7.6	5.3
Sub-total	19.3	19.5	7.5	11.2	5.5	2.9
Total	19.4		9.3		4.2	

4.2.3 Infants' physique (height, weight, seated height, and chest circumference) and skinfold thickness significantly increased with age.

- 4.2.4 Resting HR and sit-and-reach performance (flexibility) decreased significantly with age.
- 4.2.5 The performances of walking on a balance beam, continuous jump with both feet, standing long jump, 10m × 2 shuttle run, and throwing a tennis ball significantly got improved with age.
- 4.2.6 Girls had larger skinfold thickness and flexibility while boys performed better in standing long jump and throwing a tennis ball. The results mentioned reached significant difference.
- 4.2.7 Age and gender specific normative tables in quintiles and percentile ranks were generated.

4.3 Inter-correlation between Fitness Variables

- 4.3.1 In both genders, moderate-to-high correlations ($r = 0.608-0.830$) existed among infants' BMI, skinfold thickness, and chest circumference.
- 4.3.2 There was a moderate correlation ($0.400 \leq r < 0.700$) between the performances of continuous jump with both feet and 10m × 2 shuttle run in boys only. Other correlations are very small ($r < 0.400$).

4.4 Physical Activity Participation Pattern

- 4.4.1 The top three favorite physical activities of boys were “playing in the playground” (80.4%), “swimming” (49.7%), and “ball games” (44.4%). The top three favorite physical activities of girls were “playing in the playground” (80.4%), “dance” (50.7%), and “swimming” (40.6%).
- 4.4.2 More than 30% of infants received exercise training (learning class) on “cycling” (33.5%) or “swimming” (32.0%) in the past year.
- 4.4.3 Less than 20% of infants (17.7%) spent an hour or more on outdoor activities in a day. The average time spent on outdoor activities in a day could be an indicator for measuring the amount of P.A. participation. Hong Kong infants thus were not physically active enough.
- 4.4.4 Only half of infants (48.5%) participated in P.A. with their family members at least once per week in weekends or holidays. 31.8% of infants had P.A. participation with family for 1 to 2 times per month. As high as 13.6% of infants only took part in P.A. with family once several months. Even 6.1% had none in the past year. These statistics drew our attention in the society that parents may not be aware of the importance of having physical activity with their children and sacrificed the time for heavy daily workload.

4.5 Further Analysis

Relationship between Physical Fitness and Screen Time in Infants

- 4.5.1 In age group of 5–6, screen time had significant effect on resting HR, standing long jump performance, and 10m x 2 shuttle run performance ($p < 0.05$).
- 4.5.2 Infants who accumulated two hours or more of screen time in a day had higher resting HR, and poorer standing long jump and 10m x 2 shuttle run performance than those who accumulated less than two hours a day.

Relationship between the Frequency of Family Physical Activity and the Duration of Outdoor Activity in Infants

- 4.5.3 The duration of outdoor activity significantly increased with the frequency of family P.A. (boys: $p < 0.05$, girls: $p < 0.05$)

Relationship between Physical Fitness and the Duration of Outdoor Activity in Infants

- 4.5.4 The BMI of infants aged 3–4 who accumulated one hour or more of outdoor activities each day was higher than those who accumulated less than one hour at $p < 0.05$

4.6 Discussion & Recommendations

- 4.6.1 Parents and preschool organizations were recommended to perform physical fitness assessments regularly in order to understand the change of fitness level. Referral for children in need to appropriate professionals could be made as soon as possible. Fitness measurements should be comprehensive instead of only focus on one or two items. Various fitness parameters were suggested to be evaluated separately as they were mainly independent against each other
- 4.6.2 According to the criteria recommended by the International Obesity Taskforce (IOTF), as much as 19.4% and 13.5% of infants were underweight and overweight/ obese respectively. It reflected that undernourishment and over-nutrition were both common and may develop into two opposing extremes. It was necessary to deliver the information about balanced diet to parents.

- 4.6.3 One of the innate characteristics of infants is that they love to move around all day long. P.A. should be a major component in their daily life. The National Association for Sports and Physical Education (NASPE) suggested that children aged 2 years or above should accumulate at least 60 minutes of physical activities every day. In Hong Kong, less than 20% of infants (boys: 18.0%, girls: 17.4%) met the recommendation in terms of outdoor activities, reflecting insufficient P.A. Moreover, more than half of infants could not have P.A. participation with family at least once per week. The situation drew our attention. In fact, it is crucial to nurture children to have active lifestyle starting from early childhood. It was suggested that preschool organizations and parents should share the responsibilities in modifying health behaviors of infants. For example, kindergartens may incorporate minimum 20-minute physical games into their curriculum every day while parents arranged not less than 40-minute P.A. in before-school or after-school periods. Formal training should be provided to kindergarten teachers on how to teach physical movement of their students. In addition, government and related stakeholders may think about offering more workshops and fun days so as to encourage and educate parents to arrange more family-based P.A.
- 4.6.4 “Playing in playgrounds”, “swimming”, “dance”, and “ball games” were the most favorite physical activities of infants. Related stakeholders should continue to sustain the resources in these activities, so as to increase the opportunities for infants’ participation.

5. Children

5.1 Participants

- 5.1.1 Primary schools were selected through stratified randomly sampling according to their locations and modus operandi. Children aged 7–12 were invited to participate in the Study.
- 5.1.2 A total of 18 primary schools were participated, 4 from H.K. Island, 5 from Kowloon, 5 from New Territories West, and 4 from New Territories East. There were 2,803 randomly selected samples of which 2,723 were valid (97.1%).

5.1.3 The research team compared the sample sizes of four sub-groups (boys/ girls aged 7–9; boys/ girls aged 10–12) with the “Mid-2011 Provisional Population Figures” published by the C&SD so as to allow necessary weighting calculations.

5.2 Physical Fitness Level

5.2.1 The mean values of body composition and fitness parameters were shown in the table below.

Age	7–9		10–12	
Parameters	Boys	Girls	Boys	Girls
Height (<i>cm</i>)	130.68	130.05	144.71	146.18
Weight (<i>kg</i>)	30.11	28.71	40.48	39.70
BMI (<i>kg/m²</i>)	17.41	16.80	19.11	18.39
Seated height (<i>cm</i>)	70.26	69.88	75.79	77.25
Total Skinfold (Upper Arm + Calf) (<i>mm</i>)	23.50	24.45	27.10	27.55
Chest circumference (<i>cm</i>)	66.48	65.09	73.36	73.66
Resting Systolic BP (SBP, <i>mmHg</i>)	104.62	103.91	108.88	109.72
Resting Diastolic BP (DBP, <i>mmHg</i>)	63.97	64.20	66.51	67.36
Resting HR (<i>bpm</i>)	87.53	90.34	81.56	86.28
Sit-and-reach (<i>cm</i>)	1.13	3.84	-1.15	4.76
Hand Grip (<i>kg</i>)	23.21	21.30	32.88	32.80
1-min Sit-ups (<i>rep</i>)	17.07	16.02	22.09	21.53
Standing Long Jump (<i>cm</i>)	116.24	105.16	133.62	120.72
15m PACER (<i>turns</i>)	13.20	12.31	19.97	17.57

5.2.2 According to the criteria suggested by International Obesity Task Force (IOTF), 30.6% of boys and 22.8% of girls were classified as overweight or obese (see the table below). In addition, 7.8 % of boys and 9.9% of girls were categorized as underweight.

Age	Underweight		Overweight		Obese	
	Boys (%)	Girls (%)	Boys (%)	Girls (%)	Boys (%)	Girls (%)
7-9	8.0	10.1	18.2	16.6	10.0	6.2
10-12	7.6	9.8	22.4	19.7	10.2	3.2
Sub-total	7.8	9.9	20.5	18.3	10.1	4.5
Total	8.8		19.5		7.4	

5.2.3 From the table below, 4.1% of primary school students were identified with the risk of hypertension³. Although the percentage is low, caution should be raised as healthy children at young age should not have such chronic disease.

High Blood Pressure Symptom	SBP ≥ 140 mmHg or / and DBP ≥ 90 mmHg		
Age	Boys (%)	Girls (%)	Total (%)
7-9	4.2	4.5	4.3
10-12	4.1	3.7	3.9
Total	4.2	4.1	4.1

Remarks 3 Such classification was conducted according to resting SBP ≥ 140 mmHg or/and DBP ≥ 90 mmHg. In fact, BP was correlated with students' age, height, and gender. Research team also took the age-, height-, and gender-adjusted criteria recommended by "National High Blood Pressure Education Program (NHBPEP) Working Group" and "Bureau of Health Promotion, Department of Health (Taiwan)" as well as "Taiwan Society of Internal Medicine" into account. Children with BP ≥ 95 percentile value were regarded as having high BP symptom (see the table below).

Age	Meet at least 1 of the following conditions		High Blood Pressure Symptom		
	Resting SBP (mmHg)	Resting DBP (mmHg)	Boys (%)	Girls (%)	Total (%)
7-9	≥ 122	≥ 78	19.1	17.5	18.3
10-12	≥ 126	≥ 82	16.9	19.0	17.9
Total			17.8	18.3	18.1

5.2.4 Children's height, weight, BMI, seated height, skinfold thickness, chest circumference), as well as resting SBP and DBP significantly increased with age.

- 5.2.5 Resting HR decreased significantly with age.
- 5.2.6 The performances of 1-min sit-ups, standing long jump, and 15m PACER significantly got improved with age.
- 5.2.7 Girls had larger resting HR and flexibility but smaller percent body fat while boys performed better in standing long jump, and 15m PACER. The results mentioned reached significant difference.

5.3 Inter-correlation between Fitness Variables

- 5.3.1 In both genders, high correlations ($r \geq 0.700$) existed among children's BMI, skinfold thickness, and chest circumference.
- 5.3.2 In boys, a negative moderate correlation existed between skinfold thickness and 15m PACER ($r = -0.430$), reflecting that fat children may have smaller leg muscular strength and cardiovascular fitness. On the other hand, there were moderate correlations ($r = 0.404-0.479$) among 15m PACER, 1-min sit-ups, and standing long jump. Other correlations are very small ($r < 0.400$).
- 5.3.3 In girls, a moderate correlation existed between chest circumference and handgrip strength ($r = 0.451$), reflecting children with large chest cavity may have greater strength at upper limbs. In addition, there was a moderate correlation between standing long jump and 15m PACER ($r = 0.415$). Other correlations are very small ($r < 0.400$).
- 5.3.4 Age and gender specific normative tables in quintiles and percentile ranks were generated.

5.4 Health-related Lifestyle

- 5.4.1 About 80% of children (81.1%) reported that they had sufficient or very sufficient P.A. participation.

5.4.2 According to the “American Indicator”, children who accumulated at least 60 minutes moderate-or-above intensity P.A. every day in a week were classified as physically active. To facilitate the further analysis, the original categories “active” and “inactive” were subdivided into “active”, “some active”, and “sedentary” (see the table below). Only 9.5% of boys and 7.0% of girls in the study met the recommendation based on the amount of P.A. they reported. There was a difference between the classification and children’s self-evaluation. Therefore it is suggested that school-aged children may not be aware of the minimal requirement of P.A. during childhood. Education should be reinforced to alert students about the related requirement so as to promote active lifestyle in younger ages.

Category (with modification)		Number of days accumulated at least 60 minutes moderate-or-above intensity physical activity in a week	Boys (%)	Girls (%)	Total (%)
III	Active	7 days	9.5	7.0	8.3
II	Some active	1–6 days	69.9	72.2	71.0
I	Sedentary	<1 day	20.5	20.9	20.7

5.4.3 Furthermore, a classification according to the “Baseline Indicator” was shown in the table below. Children who participated in moderate-or-above intensity P.A. at least three days a week with accumulation of 30 minutes or above per day were classified as physically active. About half of children (51.8%) met the recommendation. Boys had larger proportion of being “active” than girls (54.7% vs. 48.7%). The classification result still did not compromise with children’s self-evaluation.

Category		Accumulation of moderate-or-above intensity physical activity in a week	Boys (%)	Girls (%)	Total (%)
III	Active	≥ 90 minutes	54.7	48.7	51.8
II	Some active	30–89 minutes	35.7	44.6	40.0
I	Sedentary	< 30 minutes	9.6	6.7	8.2

5.4.4 In the past year, approximately 80% of children attended sports training as an extra-curricular activity, in which “ball games” were the most popular sports with 68.1% of children chosen, followed by “swimming” (48.8%) and “track and field” (42.0%).

- 5.4.5 It was found that about 40% of children spent more than 5 hours on sedentary activities, including study and recreation, after school in their daily living.
- 5.4.6 17.9% of children slept less than 7 hours per day. On the other hand, 32.0% of children slept 9 hours or more per day.
- 5.4.7 Over 80% of children (83.5%) had an intake of at least a bowl of vegetables everyday. Also, a similar proportion of children (84.8%) had an intake of at least one fruit daily. 64.8% of children drank at least a cup of milk beverage everyday. It was found that most of the children met the recommendations on the food pyramid made by the Department of Health. However, the perception of the importance of consuming vegetables, fruits, and milk products was not studied in this study. Children may be able to meet the requirements only because their menus were set by their schools which provided lunch box or other care takers.
- 5.4.8 Around a quarter of children took high calorie food 3 times or more in a week, such as “chocolates or candies” (27.0%), “soft drinks, paper-packed drinks or canned drinks” (24.8%), and “cup noodles or instant noodles (23.5%)” (see the table below).

Foods/ Drinks	Boys (%)	Girls (%)	Total (%)
Chocolates or candies	26.0	28.1	27.0
Soft drinks, paper-packed drinks or canned drinks	27.5	21.9	24.8
Cup noodles or instant noodles	27.7	19.0	23.5
Deep fried foods (e.g. fried chicken, pork chop outlet)	22.4	14.8	18.7
Street snacks (e.g. fish balls, siu mai)	19.6	14.2	17.0
Potato chips	15.4	11.6	13.6
Yogurts	12.5	12.6	12.5
Ice-creams	11.5	11.4	11.5
Pizzas	10.7	6.8	8.8

5.5 Attitudes and Feelings toward Physical Activity

- 5.5.1 The top three reasons of poor physical health perceived were “inadequate sleep” (62.9%), “imbalance diet” (47.2%), and “too much school works” (38.8%). “Inadequate physical activity” was accounted for 38.0% .

- 5.5.2 On a scale of 1 to 4, the three most important benefits associated with P.A. participation were “P.A. helps maintain physical health” (mean rank: 3.57), “P.A. develops various physical skills” (mean rank: 3.45), and “P.A. helps control body-weight” (mean rank: 3.42).
- 5.5.3 On a scale of 1 to 4, the three attitude statements with highest level of agreement were “I like P.A.” (mean rank: 3.61), “P.A. is very interesting” (mean rank: 3.53), and “P.A. participation makes me feel happy” (mean rank: 3.52). The results reflected that interesting physical activities which brought children happiness and aroused their emotional state could increase their participation.
- 5.5.4 On a scale of 1 to 3, the top three barriers to participation of P.A. were “bad weather” (mean rank: 2.01), “too much homework” (mean rank: 1.99), and “being too tired” (mean rank: 1.86).
- 5.5.5 Among those 59.5% of children who had the experience in playing motion-sensing sports games, 81.5% of them agreed that those games has induced a greater interest in sports participation, while about half of them (50.9%) spent an hour or more per day in playing motion-sensing sports games in the past week.

5.6 Further Analysis

Relationship between Physical Fitness and Physical Activity Level (“American Indicator”) in Children

- 5.6.1 An increase in P.A. participation had its beneficial effect on muscular strength and cardiovascular fitness at significant level. Related test items included handgrip, 1-min sit-ups, standing long jump, and 15m PACER.
- 5.6.2 “Active” boys demonstrated 8.4% and 19.8% better performance in standing long jump and 15m PACER respectively than “sedentary” boys who also demonstrated 9.0% and 14.7% lower performance in handgrip and 1-min sit-up than “some-active” boys.
- 5.6.3 In girls, main difference mainly existed between “some-active” children and “sedentary” children.

Relationship between Screen Time and Physical Activity Level (“American Indicator”) in Children

- 5.6.4 When children’s screen time (watching TV/ movies, playing electronic games, or using computer/ web-surfing) increased, their P.A. level significantly dropped.

- 5.6.5 In boys, there were a higher proportion of children who spent more than 3 hours on screen activities during school days classified as “sedentary” than those with 3-hour-or-less screen time (32.0% vs. 18.1%).
- 5.6.6 In girls, there were fewer children who spent an hour or above on screen activities classified as “active” than those with less than an hour screen time (5.3% vs. 9.7%).

Relationship between Screen Time and Body Mass Index in Children

- 5.6.7 When children’s screen time increased, their BMI significantly rose.
- 5.6.8 Boys who spent 2 hours or above on screen activities during school days had greater BMI than those without screen time.
- 5.6.9 Girls who spent more than 3 hours on screen activities had greater BMI than those with less than 1 hour or no screen time.

Relationship between Physical Fitness and Sleeping Time in Children

- 5.6.10 There was a significant difference in fitness level between groups of children having different sleep times.
- 5.6.11 Children who slept for 7 to 8 hours daily performed better in standing long jump and 15m PACER than those who slept less or more.
- 5.6.12 In 1-min sit-ups test, boys with 7 to 8 sleeping hours daily also had better performance whereas girls with 9 sleeping hours or more showed worse performance than those who slept less.
- 5.6.13 In handgrip test, children who slept 9 hours or more demonstrated smaller forearm strength.

Relationship between Parent’s Exercise Frequency and Children’s Physical Activity Level (“American Indicator”)

- 5.6.14 There were significant differences in children’s P.A. levels between groups of parents with various exercise frequencies per week. Children tended to be more physically active if their fathers or mothers were more physically active.
- 5.6.15 When fathers did exercise at least 6 times per week, 24.1% of their sons and 16.2% of daughters were “active”. The proportions were higher than those whose fathers had no exercise in a week (boys: 7.5%, girls: 4.9%).

5.6.16 When mothers' exercise frequency reached 6 times per week, 23.6% of their sons and 15.0% of daughters were "active". The proportions were also higher than those whose mothers had no exercise in a week (boys: 6.5%, girls: 5.4%).

5.6.17 In fact, there were 34.9% of fathers and 37.2% of mothers who did not participate in any form of exercise. Moreover, as much as 62.9% of family did not manage to have regular family activities in sports weekly during weekends or holidays in the past year.

Comparison of Physical Activity Level across Years

5.6.18 The results from the table below showed the comparison of P.A. level based on the "Baseline Indicator" between "Sports for All Study 2009" and the current study using the Chi-square test. There was no significant difference in P.A. level between the two years.

Study	P.A. Level		Chi-square Test <i>p</i> -value
	Some active, Sedentary (%)	Active (%)	
2009	50.8	49.2	0.147
2011	47.7	52.3	

5.6.19 Chi-square test was also used to compare the P.A. level based on "American Indicator" proposed by USDHHS between "Sports for All Study 2009" and the current study (see the table below). More children (9.0%) met the guidelines requirement in the current study, higher than the proportion reported in Sports for All Survey (5.7%). The difference reached statistical significance. Such results were mainly attributed to a significant increasing percentage of girls being classified as physically active from 3.7% to 7.7%.

Study	P.A. Level		Chi-square Test <i>p</i> -value
	Some active, Sedentary (%)	Active (%)	
2009	94.3	5.7	0.006 [@]
2011	91.0	9.0	

[@] Significant difference at $p \leq 0.05$

5.7 Discussion & Recommendations

- 5.7.1 To allow children to have full understanding of how much P.A. is recommended for their age, corresponding frequency, duration, and intensity under the two P.A. indicators should be explained clearly to them.
- 5.7.2 According to the criteria suggested by the International Obesity Taskforce (IOTF), 19.5% and 7.4% of children were overweight and obese respectively. Other than insufficient P.A. participation, children may suffer from over-nutrition. Related stakeholders are advised to publicize the impact of eating junk food (generally referring to fast food and low nutritional value food) on health.
- 5.7.3 To alleviate the main constraint “bad weather” to P.A. participation, it is suggested that related stakeholders establish more highly-accessible indoor activity facilities and play rooms suitable for children aged 7 to 12.
- 5.7.4 To awaken children’s interests in do exercise and raise their P.A. levels, schools and families are recommended to arrange diversified exercise games in groups so as to provide children with interesting and delighted experiences.
- 5.7.5 Parents’ and children’s P.A. levels were found to be highly correlated. Parents’ positive attitude toward P.A. participation had great influence on children. It is suggested to let parents obtain more information about the benefits of developing regular exercise habit and encourage them to organize family-based P.A. functions.
- 5.7.6 Screen-based motion-sensing sport games become more popular nowadays. They involve intensive body movements that simulate biomechanics of various sport activities in reality. Motion-sensing sports games are thought to be one of the innovative and attractive means which could be supplementary to traditional mode of exercise promotion. Certainly, more evidences should be explored to evaluate their effectiveness. Do playing motion-sensing sports games increase overall fitness level? Are cognitive and psychosocial effects brought by playing motion-sensing sports games identical to those of actual P.A. or sports participation? Are any safety measures such warm-up and cool-down activity incorporated in the motion-sensing games incorporated?

6. Adolescents

6.1 Participants

- 6.1.1 One secondary school was selected from 18 districts respectively through stratified randomly sampling. Adolescents aged 13–19 were invited to participate in the Study.
- 6.1.2 There were 2,805 randomly selected samples of which 2,517 were valid (89.7%).
- 6.1.3 The research team compared the sample sizes of four sub-groups (boys/ girls aged 13–15; boys/ girls aged 16–19) with the “Mid-2011 Provisional Population Figures” published by the C&SD so as to allow necessary weighting calculations.

6.2 Physical Fitness Level

- 6.2.1 The mean values of body composition and fitness parameters were shown in the table below.

Age	13–15		16–19	
Parameters	Boys	Girls	Boys	Girls
Height (<i>cm</i>)	165.46	157.26	170.99	158.90
Weight (<i>kg</i>)	55.83	49.56	62.30	51.06
BMI (<i>kg/m²</i>)	20.32	20.02	21.27	20.19
Seated height (<i>cm</i>)	86.75	83.98	90.76	85.12
Chest circumference (<i>cm</i>)	82.05	80.63	86.08	81.88
Total Skinfold (Upper Arm + Calf) (<i>mm</i>)	21.88	28.95	19.18	28.97
Percent Body Fat	18.23	24.08	14.40	22.93
Resting Systolic BP (SBP, <i>mmHg</i>)	115.46	107.20	122.48	107.28
Resting Diastolic BP (DBP, <i>mmHg</i>)	69.11	66.70	73.56	66.70
Resting HR (<i>bpm</i>)	81.24	84.91	76.15	83.07
Sit-and-reach (<i>cm</i>)	-0.43	6.84	2.48	7.81
Hand Grip (<i>kg</i>)	58.37	40.65	71.97	42.43
1-min Sit-ups (<i>rep</i>)	28.61	22.03	30.62	22.58
Push-ups (<i>rep</i>)	9.51	6.19	15.06	6.68
Standing Long Jump (<i>cm</i>)	169.44	131.31	187.75	134.93
15m PACER (<i>turns</i>)	38.05	21.77	43.50	22.23

6.2.2 According to the criteria suggested by International Obesity Task Force (IOTF), 18.5% of boys and 9.5% of girls were classified as overweight or obese (see the table below). In addition, 15.5 % of boys and 21.0% of girls were categorized as underweight.

Age	Underweight		Overweight		Obese	
	Boys (%)	Girls (%)	Boys (%)	Girls (%)	Boys (%)	Girls (%)
13–15	12.8	13.4	15.0	9.5	7.1	2.4
16–19	17.3	26.3	11.9	5.8	4.0	1.8
Sub-total	15.5	21.0	13.2	7.4	5.3	2.1
Total	18.2		10.3		3.7	

6.2.3 According to SBP $\geq 140\text{mmHg}$ or / and DBP $\geq 90\text{mmHg}$, 7.5% of boys and 1.2% of girls were identified with the risk of hypertension ⁴ (see the table below).

High Blood Pressure Symptom	SBP $\geq 140\text{mmHg}$ or / and DBP $\geq 90\text{mmHg}$		
Age	Boys (%)	Girls (%)	Total (%)
13–15	3.4	1.4	2.4
16–19	10.3	1.1	5.8
Total	7.5	1.2	4.4

Remarks 4

Such classification was conducted according to resting SBP $\geq 140\text{ mmHg}$ or/and DBP $\geq 90\text{ mmHg}$. In fact, BP was correlated with students' age, height, and gender. Research team also took the age-, height-, and gender-adjusted criteria recommended by “National High Blood Pressure Education Program (NHBPEP) Working Group” and “Bureau of Health Promotion, Department of Health (Taiwan)” as well as “Taiwan Society of Internal Medicine” into account. Adolescents with BP ≥ 95 percentile value were regarded as having high BP symptom (see the table below).

Age	Meet at least 1 of the following conditions		High Blood Pressure Symptom		
	Resting SBP (<i>mmHg</i>)	Resting DBP (<i>mmHg</i>)	Boys (%)	Girls (%)	Total (%)
13–15	≥ 136	≥ 86	8.3	2.4	5.4
16–19	16–18	≥ 142	7.5	0.6	4.1
	19	≥ 140			
Total			7.8	1.3	4.6

- 6.2.4 The exercise performances of handgrip, 1-min sit-ups, push-ups, standing long jump, and 15m PACER significantly got improved with age.
- 6.2.5 Boys had greater height, weight, seated height, chest circumference, and resting SBP and DBP than girls while girls possessed higher skinfold thickness, percent body fat, and resting HR than boys. The results mentioned reached significant difference.

6.3 Inter-correlation between Fitness Variables

- 6.3.1 In both genders, moderate-to-high correlations ($r = 0.625\text{--}0.897$) existed among adolescents' BMI, skinfold thickness, and chest circumference. There were also moderate correlations between resting SBP and BMI (boys: $r = 0.478$, girls: $r = 0.423$) and chest circumference (boys: $r = 0.467$, girls: $r = 0.386$).
- 6.3.2 In boys, moderate correlations were found between body compositions and exercise performances (chest and handgrip: $r = 0.402$, skinfold thickness and 15m PACER: $r = -0.408$, and skinfold thickness and standing long jump: $r = -0.422$). Other correlations are very small ($r < 0.400$).
- 6.3.3 On the other hand, the exercise performances of 1-min sit-ups, push-ups, 15m PACER, and standing long jump in boys were moderately inter-correlated ($0.400 \leq r < 0.700$) except for the associations between 1-min sit-ups and 15m PACER ($r = 0.341$) and standing long jump ($r = 0.398$). In girls, a moderate correlation was shown only between 15m PACER and standing long jump ($r = 0.471$). No other moderate-or-above inter-correlations were found.
- 6.3.4 Age and gender specific normative tables in quintiles and percentile ranks were generated.

6.4 Health-related Lifestyle

6.4.1 According to the “American Indicator”, adolescents who accumulated at least 60 minutes moderate-or-above intensity P.A. every day in a week in which any three days involved with vigorous intensity P.A. were classified as physically active. To facilitate the further analysis, the original categories “active” and “inactive” were subdivided into “active”, “some active”, and “sedentary” (see the table below). Only 12.8% of boys and 4.2% of girls in the study met the recommendation based on the amount of physical activity they reported.

Category (with modification)		P.A. Level	Boys (%)	Girls (%)	Total (%)
III	Active	Accumulation of at least 60 minutes moderate-or-above intensity physical activity everyday in a week in which any three days involved with vigorous intensity physical activity	12.8	4.2	8.4
II	Some active	Accumulation of at least 10 minutes but less than 60 minutes moderate-or-above intensity physical activity everyday in a week	47.6	41.9	44.7
I	Sedentary	Accumulation of less than 10 minutes moderate-or-above intensity physical activity everyday in a week	39.6	53.9	46.9

6.4.2 On the other hand, a classification according to the “Baseline Indicator” was shown in the table below. Adolescents who participated in moderate-or-above intensity P.A. at least three days a week with accumulation of 30 minutes or above per day were classified as physically active. 42.3% of adolescents were “active” whereas 28.9% of adolescents were “sedentary”, 20.7 percent points higher than the prevalence in children (8.2%). 50.1% of boys were “active”, 15.9 percent points higher than the prevalence in girls.

Category		Accumulation of moderate-or-above intensity physical activity in a week	Boys (%)	Girls (%)	Total (%)
III	Active	≥ 90 minutes	50.1	34.2	42.3
II	Some active	30–89 minutes	25.3	32.5	28.8
I	Sedentary	< 30 minutes	24.6	33.4	28.9

- 6.4.3 The top three extra-curricular activities during weekend were “friends gathering” (61.5%), “sports” (40.5%), and “visual/ audio entertainment” (35.0%). In girls, the top three activities were “friends gathering” (64.0%), “shopping” (48.1%), and “visual/ audio entertainment” (44.6%).
- 6.4.4 A majority of adolescents (boys: 62.6%, girls: 60.4%) felt that homework and studying was generally acceptable. Almost one-third of adolescents (boys: 30.3%, girls: 36.7%) felt very hard.
- 6.4.5 During school days, a majority of adolescents (boys: 81.0%, girls: 80.2%) slept at least 6 but less than 9 hours. 8.8% of boys and 14.4% of girls slept less than 6 hours whereas 10.2% of boys and 5.4% of girls slept over 9 hours.
- 6.4.6 Senior secondary students (boys: 7.12 *hours*, girls: 6.74 *hours*) slept less than junior secondary students (boys: 7.64 *hours*, girls: 7.32 *hours*), which may reflect academic pressure in senior secondary classes. On the other hand, girls slept less than boys during school days but slept more than boys during non-school days
- 6.4.7 25.7% of adolescents spent 3 hours or more on screen activities (including watching TV, using cell phone, computer, and video games) for an average day.

6.5 Physical Activity Habits and Altitude & Preferences towards Physical Activities

- 6.5.1 The three most popular sports activities were “ball games” (75.1%), “ice skating or roller skating” (24.8%), and “swimming” (24.7%).
- 6.5.2 The top three reasons adolescents selected for poor physical conditions were “inadequate sleep” (75.5%), “inadequate sports training” (55.8%), and “stress” (34.2%).
- 6.5.3 There were 56.9% of boys and 28.3% of girls who really liked to attend P.E. lessons. Only 5.0% and 10.9% of boys and girls respectively disliked the lessons. The rest of them included “fairly liked” and “no comment”.
- 6.5.4 68.3% of adolescents said that they would participate or actively participate in sports activities.

- 6.5.5 The top three reasons for participating in sports activities were “raising ability on sports” (55.3%), “health strengthening/ prevention or cure of sickness” (51.3%), and “releasing pressure and emotions” (38.7%). “Keeping fit” was the fourth popular reason (36.3%) in girls, 15.4 percent points higher than boys.
- 6.5.6 The top three barriers to sports activities participation were “not interested” (52.3%), “lazy” (45.2%), and “lack of spare time” (44.3%).
- 6.5.7 The top three reasons for participating in distance run training were “raising ability on sports” (55.8%), “health strengthening/ prevention or cure of sickness” (47.3%), and “releasing pressure and emotions” (27.0%). The distribution was similar to the reasons for participating in sports activities.
- 6.5.8 The top three barriers to distance run training participation were “tired” (58.8%), “not interested” (55.0%), and “lazy” (35.7%).
- 6.5.9 The top three reasons for participating in strength training were “raising ability on sports” (53.1%), “Keeping fit” (48.3%), and “health strengthening/ preventing or cure of sickness” (47.3%).
- 6.5.10 The major reasons for not participating in strength training were “not interested” (52.4%), “tired” (46.2%), and “lazy” (35.3%). The major reasons were the same as not participating in distance run.
- 6.5.11 40.8% of boys’ parents and 30.6% of girls’ parents always or sometimes recommended their children to reduce sport activities for studies.

6.6 Further Analysis

Main Effects and Interaction Effect of Gender and Age-Groups on Physical Fitness

- 6.6.1 In 2-way ANOVA, significant main effects (age-groups and gender) and interaction effect (gender × age-groups) were found on physical fitness. For exercise measurements, boys performed better in handgrip, 1-min sit-ups, push-ups, standing long jump, and 15m PACER than girls except for the sit-and-reach test.
- 6.6.2 The overall fitness level significantly increased with age in both adolescents in which boys showed a greater improvement than girls.

Relationship between Physical Fitness and Physical Activity Levels (“American Indicator”) in Adolescents

- 6.6.3 P.A. levels did not cause significant effects on BMI, percent body fat, and chest circumference in both genders. It may reflect the fact that there are large variations of growth factors between individuals at puberty stage.
- 6.6.4 Nevertheless, P.A. levels seemed to have significantly beneficial effects of decreasing resting HR as well as enhancing other exercise performances (sit-and-reach, 1-min sit-ups, push-ups, handgrip, standing long jump, and 15m PACER). Similar findings were found for both boys and girls.

Influence of Parental Education Level on Fitness Parameters in Adolescents

- 6.6.5 In fathers with tertiary education attainment, their sons had smaller chest circumference and forearm strength but lower BMI and greater cardiovascular fitness in terms of better performance in 15m PACER test. Girls whose fathers attained tertiary education level possessed higher abdominal endurance and cardiovascular fitness.
- 6.6.6 In mothers with primary-or-below education attainment, their sons had greater flexibility, forearm strength, and pectoral muscular endurance. There was no significant difference found in fitness parameters among various mother’s education levels in girls.

Relationship between Parents’ and Adolescents’ Physical Activity Levels

- 6.6.7 Significant differences were found between adolescents’ and fathers’ P.A. levels. The P.A. levels of boys and girls with active father (frequency of exercise training ≥ 3 per week) would have respectively about 38% and 96% higher in terms of MET-minute than those with sedentary father (frequency of exercise training < 1). Active mothers were found to be associated with more physical activities in girls.
- 6.6.8 In other words, adolescents tended to be more active if the parents were more active. However, 47.5% of fathers and 52.1% of mothers did not take part in exercise training in the past year. Moreover, as high as 77.0% of adolescents did not participate in P.A. with their families regularly (at least once per month).

Relationship between Screen Time and Physical Fitness in Adolescents

6.6.9 Adolescents were grouped according to the amount of screen time. Boys who spent 3 hours or more on screen activities had poorer flexibility and cardiovascular fitness than those having less than 3 hours screen activity. Furthermore, they also were found to have lower muscular endurance (abdomen and pectoris) and power (lower limbs) than those with less than 2 hours on screen activities. There was no significant difference in fitness parameters among various amount of screen time in girls.

Comparison of Physical Activity Level across Years

6.6.10 The results from the table below showed the comparison of P.A. level based on the “Baseline Indicator” between “Sports for All Study 2009” and the current study. The proportion of “active” adolescents dropped significantly from 48.6% to 42.1% in these two years. Such results were mainly attributed to a significant decreasing percentage of boys being classified as physically active from 57.5% to 50.1%.

Study	P.A. Level		Chi-square Test <i>p</i> -value
	Some active, Sedentary (%)	Active (%)	
2009	51.4	48.6	0.001 [@]
2011	57.9	42.1	

[@] Significant difference at $p \leq 0.05$

6.6.11 The comparison analysis of P.A. levels based on the “American Indicator” was not available as the data collected from the two studies could not be matched.

Preliminary Study on the “Sport For All Day”

6.6.12 Close to half of adolescents (45.3%) adolescent heard about the “Sport for All Day” organized by the LCSD to provide Hong Kong people with a series of free recreation and sports programmes in the past two years (2010, 2011). No significant difference was observed between genders as shown in the table below.

Heard about the “Sport For All Day” Before	Boys (%)	Girls (%)	Total (%)	Chi-square Test <i>p</i> -value
Yes	44.7	45.7	45.2	0.607
No/ Not sure	55.3	54.3	54.8	

6.6.13 32.3% of the respondents who were aware of the “Sport for All Day” participated in the event. There was a significant difference between genders that more boys took part in the event, 7.2 percent points higher than girls as shown in the table below.

Participation in the “Sport For All Day”	Boys (%)	Girls (%)	Total (%)	Chi-square Test <i>p</i> -value
Yes	35.8	28.6	32.3	0.009 @
No/ Not sure	64.2	71.4	67.7	

@ significant difference at $p \leq 0.05$

6.6.14 As much as 67.4% of “Sport for All Day” participants thought that their interests in sports and P.A. could be raised through the participation. A significant difference was found between genders as shown in the table below. More boys (71.8%) had such thought than girls (61.9%).

Increase Interests in Sports and P.A.	Boys (%)	Girls (%)	Total (%)	Chi-square Test <i>p</i> -value
Yes	71.8	61.9	67.4	0.042 @
No	28.2	38.1	32.6	

@ significant difference at $p \leq 0.05$

6.7 Discussion & Recommendations

6.7.1 In the current study, a majority of the physical fitness parameters showed significant difference between age groups of 13–15 and 16–19. Gender difference was also found in both age groups. This reflected the growth spurt in physical development during puberty. Physical education curriculums (e.g. exercise intensity, equipment disposal, and safety measures) in schools are recommended to cater for the growth needs. Schools were also suggested to take reference from “Physical Education Key Learning Area Curriculum Guide (2002)” published by the EDB so as to provide adolescents with diversified learning experiences through a series of physical activities.

- 6.7.2 According to the criteria suggested by the International Obesity Taskforce (IOTF), 18.5% of boys and 9.4% of girls were overweight or obese. On the other extreme, 15.5% of boys and 21.0% of girls were underweight. Boys tended to encounter overweight problem while girls tended to have underweight problem. One possible reason was the promotion of “Thinness is beauty” in girls by mass media. It was suggested that related stakeholders proactively publicize the correct concepts of achieving a healthy body weight as well as how to gain/lose weight in a proper way. Adolescents were encouraged to keep track on their own weight and percent body fat.
- 6.7.3 Findings revealed that more than 30% of parents (35.8%) frequently or occasionally recommended their children to reduce sport activities for studies. It is believed that adolescents may treat studies as a higher priority so as to satisfy the traditional expectations of parents and social norms. The stress from studying was not that little in adolescents. An increase in study stress may be one of the reasons for the decrease in P.A. participation. Therefore related stakeholders should work together to find solutions in order to mitigate the problems and promote healthier lifestyle other than academic pursuit alone. The benefits of engaging in P.A. should not only be promoted to students, but should also be emphasized to their parents who would be the role model to show more active lifestyle to their children. Family activities in sports and exercises were encouraged.
- 6.7.4 With reference to the key barrier “not interested” to sports activities participation, great variety of sports content, such as cycling, sailing, canoeing, and bowling, etc., should be introduced to adolescents. “Ball games”, “ice skating or roller skating”, and “swimming” may also be continuously promoted with corresponding input of resources as they were reported to be the most popular sports.
- 6.7.5 As can be seen from the survey results, 62.8% of adolescents chose “friends gathering” as their major extra-curricular activities during weekend. “No company” was the fifth place of the reasons why adolescents did not participate in sports activities. Close to 20% of adolescents (19.7%) regarded it as a key factor. It is possible that if the norms of peer group prefer sedentary activities, active adolescents are likely to conform to these peer norms and become less active, vice versa. Peer group intervention is therefore suggested to be one of the effective strategies to promote physical activity in adolescents.

7. Adults & Elderly

7.1 Participants

- 7.1.1 After a try-out test for Hong Kong residents aged 20 to 69 was implemented, a Territory-wide Physical Fitness Test for the same target group was conducted in 18 districts, from April 2011 to January 2012. To begin with, Hong Kong residents were stratified randomly selected for a face-to-face interview, then respondents were invited to attend fitness assessments in a sports center that near their home by appointment. To complete all the field fitness testing, fifty-seven different sports centers provided by the LCSD and 270 testing sessions were utilized.
- 7.1.2 7,591 households accepted the interview which involved 15,013 household members. There were 2,354 sets of valid data (15.7%) who completed both the interview and physical fitness tests.
- 7.1.3 Although the sampling percentages of middle-aged adults and elderly were close to the actual population distribution ratio, young adults contributed a relatively small portion for both genders and diverged from the sampling requirement. The research team compared the sample sizes of six sub-groups (men/ women aged 20–39; men/ women aged 40–59; men/ women aged 60–69) with the “Mid-2011 Provisional Population Figures” published by the C&SD so as to allow necessary weighting calculations.

7.2 Physical Fitness Level

7.2.1 The mean values of body composition and fitness parameters were shown in the table below.

Age		20–39		40–59		60–69	
Parameters	Gender	Men	Women	Men	Women	Men	Women
Body Height (<i>cm</i>)		171.32	158.25	167.91	156.10	165.46	153.81
Body Weight (<i>kg</i>)		69.74	53.70	68.91	56.92	65.24	57.78
BMI (<i>kg/m²</i>)		23.72	21.44	24.41	23.35	23.81	24.43
Chest Circumference (<i>cm</i>)		91.85	83.32	92.98	87.31	91.37	89.12
Waist Circumference (<i>cm</i>)		82.97	71.18	86.36	77.33	86.83	82.18
Hip Circumference (<i>cm</i>)		95.15	91.52	94.64	94.09	93.05	95.72
Skinfold Thickness	Upper Arm (<i>mm</i>)	11.35	16.43	9.95	17.68	9.45	17.65
	Subscapular (<i>mm</i>)	16.79	16.71	18.40	20.47	17.13	20.34
	Abdomen (<i>mm</i>)	21.31	20.08	22.18	22.89	20.53	24.29
	Chest (<i>mm</i>)	11.77	-----	13.58	-----	13.29	-----
	Suprailiac (<i>mm</i>)	-----	16.03	-----	18.85	-----	19.54
	Thigh (<i>mm</i>)	13.28	23.20	12.01	23.82	10.95	21.38
Resting HR (<i>bpm</i>)		74.85	75.63	74.29	74.68	73.75	72.51
Resting Systolic BP (<i>mmHg</i>)		124.31	110.54	129.16	123.08	135.02	132.63
Resting Diastolic BP (<i>mmHg</i>)		78.05	70.06	81.89	75.40	80.37	75.49
3-min Step Test (Post Exercise HR)		143.85	145.91	132.93	144.07	-----	-----
3-min Step Test (Recovery HR)		122.13	121.77	113.96	119.54	-----	-----
Sit-and-reach (<i>cm</i>)		-3.01	2.81	-2.58	3.22	-4.80	1.81
Hand Grip (<i>kg</i>)		77.94	45.56	74.47	43.46	66.15	38.77
1-min Sit-up (<i>rep</i>)		25.00	17.99	-----	-----	-----	-----
Push-up (<i>rep</i>)		17.07	9.17	-----	-----	-----	-----
Vertical Jump (<i>cm</i>)		36.16	22.74	-----	-----	-----	-----
Closed Eyes Balance Stand (<i>s</i>)		39.31	32.23	15.53	14.42	9.80	8.76
Reaction Time (<i>s</i>)		0.4076	0.4623	0.4504	0.5033	0.5134	0.5543

7.2.2 According to BMI classification, 14.0% of young women were regarded as underweight (see the table below). The proportions of overweight and obese individuals increased in the middle-aged. The prevalence of women experienced the obesity problem kept on rising. 40.3% of men aged 40–59 and 40.8% of women aged 60–69 were obese.

Age	Underweight (%)		Normal (%)		Overweight (%)		Obese (%)	
	BMI < 18.5		18.5 ≤ BMI < 23		23 ≤ BMI < 25		BMI ≥ 25	
	Men	Women	Men	Women	Men	Women	Men	Women
20–39	6.8	14.0	39.1	60.1	19.3	13.3	34.8	12.6
40–59	2.1	3.5	31.4	48.1	26.2	21.7	40.3	26.7
60–69	5.4	3.0	31.7	37.1	28.7	19.1	34.2	40.8
Sub-total	4.3	7.9	34.4	52.0	23.9	17.8	37.3	22.3
Total	6.3		44.0		20.6		29.1	

7.2.3 Men and women with waist circumference greater than 90cm and 80cm respectively were classified as having central obesity. Over a quarter of adults (men: 28.1%, women: 25.9%) encountered related issue as shown in the table below.

Waist Circumference	Central Obesity (%)		
	> 90cm	> 80cm	Total
Age	Men	Women	
20–39	22.2	10.8	15.7
40–59	31.4	32.7	32.1
60–69	33.1	55.1	44.0
All	28.1	25.9	26.9

7.2.4 There was a significant increase in prevalence of hypertensive response from young to old adulthood as shown in the table below. 16.4% and 6.3% of young men and women suffered from the risks of hypertension. The percentage of older men with high blood pressure symptom was approximately 2.4 times the young men. A 5.6-time expansion even existed in women.

Age	High Blood Pressure Symptom (%)		
	Men	Women	Total
20–39	16.4	6.3	10.6
40–59	29.0	19.4	23.9
60–69	40.0	35.1	37.6
All	25.7	15.6	20.2

7.3 Inter-correlation between Fitness Variables

7.3.1 Among 14 fitness parameters, high correlations ($r \geq 0.700$) existed between BMI and skinfold thickness (except for older women), resting SBP and DBP (except for the elderly), and step test post-exercise HR and recovery HR (except for young women). Moderate correlations ($0.400 \leq r < 0.700$) were only found among a few parameters. Other correlations are very small ($r < 0.400$).

7.3.2 The overall result suggests that most fitness parameters are quite unique and independent to each other. Hence, when evaluating physical fitness, various aspects have to be included for a completed fitness evaluation.

7.3.3 Age and gender specific normative tables in quintiles and percentile ranks were generated.

7.4 Health-related Lifestyle

7.4.1 The research team took reference from the International Physical Activity Questionnaire (IPAQ) to help transform the time spent on moderate-or-above intensity P.A. into MET-minutes. According to the “American Indicator”, adults and elderly who accumulated at least 150 minutes moderate-or-above intensity physical activity in a week (or 75minutes vigorous intensity P.A. or other combinations of different intensities, e.g. accumulation of 90 minutes and 30 minutes moderate- and vigorous- intensity P.A. respectively) were classified as physically active. To facilitate the further analysis, the original categories “active” and “inactive” were subdivided into “highly active”, “active”, “some active”, and “sedentary” (see the table below). Less than 30% of adults were categorized as having sufficient P.A. (highly active and active levels) in all age groups. For almost every two adults, one of them was found sedentary.

Category (with modification)		Accumulation of moderate-or-above intensity physical activity in a week	Age (%)			Total (%)
			20–39	40–59	60–69	
IV	Highly active	≥ 300 minutes	14.7	16.3	16.7	15.7
III	Active	150–299 minutes	13.8	12.2	12.6	12.9
II	Some active	31–149 minutes	26.4	20.5	18.6	22.7
I	Sedentary	≤ 30 minutes	45.1	51.0	52.1	48.7

7.4.2 On the other hand, a classification according to the “Baseline Indicator” was shown in the table below. Adults and elderly who participated in moderate-or-above intensity P.A. at least three days a week with accumulation of 30 minutes or above per day were classified as physically active. 37.1% of adults and elderly were “active”. 41.3% of men were “active”, 7.7 percent points higher than the prevalence in women.

Category		Accumulation of moderate-or-above intensity physical activity in a week	Boys (%)	Girls (%)	Total (%)
III	Active	≥ 90 minutes	41.3	33.6	37.1
II	Some active	30–89 minutes	16.8	19.6	18.3
I	Sedentary	< 30 minutes	41.9	46.8	44.6

- 7.4.3 The majority of adults and elderly (71.1%) slept 7 hours or more but less than 9 hours per day on working or school days. There were 5.5% sleeping less than 6 hours per day. People slept about 7.46 hours on average.
- 7.4.4 The top four activities engaged in leisure time were “audio/ visual entertainment” (63.0%), “shopping” (45.2%), “browsing on the internet” (40.2%), and “playing sports or doing physical exercise” (27.6%).
- 7.4.5 In adults and elderly, the average walking time per day was 84.90 minutes. More than 60% of them (62%) walked less than 120 minutes. On the other hand, the average sitting time per day was 7.01 hours.
- 7.4.6 More than 50% of samples (52.3%) took part in sports activities for “health strengthening/ prevention or cure of sickness”. As high as a quarter of young adults (27.5%) did not have any involvement in sports activities. The prevalence of other two age groups got even higher (middle-aged: 35.7%, elderly: 32.5%). The situation raised awareness. More and more people lost interests in playing sports as getting older. The percentage of unsuitability to participate for health concern also increased with age.
- 7.4.7 The top three barriers to sports activities participation were “lack of spare time” (62.9%), “tired”(37.9%), and “lazy” (33.5%). 71.5% of young adults expressed that they did not have spare time to take part in sports activities. This barrier contributed a relatively smaller proportion to other age groups (middle-aged: 61.2%, elderly: 40.8%).
- 7.4.8 The most participated sports activities were “ball games” (40.4%) and “running/ jogging” (36.9%) in young adults. “Walking” become more popular and got the first place in older adults (51.4%). As age increased, physically demanding activities or those involving body-contact lost popularity. Walking including hiking was more preferable for the elderly.

7.5 Further Analysis

Main Effects and Interaction Effect of Gender and Age-Groups on Physical Fitness

- 7.5.1 In 2-way ANOVA, significant main effects (age-groups and gender) and interaction effect (gender \times age-groups) were found on physical fitness. For exercise measurements, men performed better in handgrip, 1-min sit-ups, push-ups, vertical jump, closed eyes balance stand, reaction time, and 3-min step test than women except for the sit-and-reach test. The overall fitness level significantly decreased with age in both genders.
- 7.5.2 These results suggest that women decline in fitness parameters (including resting SBP and DBP, BMI, waist circumference, hip circumference, percent body fat, handgrip, estimated VO₂ max, and step test recovery HR in 3-min step test) much quicker than men from young adult to middle aged and carry over to older adult years. More attention was suggested to promote health-related fitness of women.

Relationship between Physical Fitness and Physical Activity Level (“American Indicator”) in Adults & Elderly

- 7.5.3 Significant differences in many physical fitness parameters were found amongst the three P.A. levels in both genders. Men with “sufficient P.A.” (active and highly active) possessed lower resting HR, waist-to-hip ratio, higher cardiovascular fitness, flexibility, forearm strength, abdominal endurance, and shorter reaction time, but larger BMI. In addition, men having “sufficient” or “some P.A.” (some active) both had greater pectoral endurance and leg strength.
- 7.5.4 Sedentary group generally possessed lower physical fitness level.
- 7.5.5 Women had similar findings as men.

Relationship between Family Income and Physical Activity Level (“American Indicator”)

- 7.5.6 Family income was found to be highly correlated with the P.A. level. 40.0% of men whose family income was higher than the median income were “active”. Such prevalence was higher than those below the median income (28.7%). In other words, more physically active men were found in family with higher income.
- 7.5.7 On the contrary, family income did not indicate a significant relationship with the P.A level of women.

Relationship between Smoking Status and Physical Fitness

- 7.5.8 Significant differences were found in several physical fitness parameters between smokers and non-smokers in men. The sampling error may be high as the number of female smoking participants was so small. Therefore significant difference in fitness profiles could not be easily defined between female smokers and non-smokers.
- 7.5.9 In men, non-smokers possessed smaller waist-to-hip ratio, larger leg power, better balance-controlling ability, and shorter reaction time.

Relationship between Occupations and Physical Activity Level (“American Indicator”)

- 7.5.10 Types of occupations were highly correlated with employee’s P.A. level.
- 7.5.11 The occupation with highest proportion of “active” level was professionals (47.9%) in men and managers or administrators (34.8%) in women.

Relationship between Physical Fitness and Educational Attainment

- 7.5.12 Significant differences were found in many physical fitness parameters among various groups of educational attainment in both genders. Women with tertiary education level had smaller skinfold thickness, better cardio-respiratory fitness (in terms of estimated VO₂ max), and larger abdominal endurance and leg strength. Moreover, women with higher levels of education had smaller BMI, percent body fat, chest circumference, waist circumference, hip circumference, waist-to-hip ratio, resting SBP and DBP, greater handgrip strength and balance-controlling ability, and shorter reaction time. More desirable fitness level may be attributed to their consciousness of health and fitness.
- 7.5.13 Men had similar findings but not such obviously among different education levels existed in women. Significant differences were involved in some fitness parameters mainly between secondary level and tertiary level of education. That is, men with tertiary education attainment were found to better fitness level (including lower BMI, percent body fat, and resting DBP, as well as greater muscular strength at abdomen and lower limbs) than those having secondary level of education.

Comparison of Physical Activity Level across Years

7.5.14 The results from the table below showed the comparison of P.A. level based on the “Baseline Indicator” between “Sports for All Study 2009” and the current study. The proportion of “active” adults and elderly dropped significantly from 43.5% to 37.0% in these two years. Such results were attributed to a significant decrease in the prevalence of men (↓8.7 percent points) and women (↓4.2 percent points) being classified as physically active.

Study	P.A. Level		Chi-square Test <i>p</i> -value
	Some active, Sedentary (%)	Active (%)	
2009	56.5	43.5	0.000 [@]
2011	63.0	37.0	

[@] Significant difference at $p \leq 0.05$

7.5.15 On the other hand, the P.A. level based on “American Indicator” classification was compared between “Sports for All Study 2009” and the current study (see the table below). There was no significant difference in P.A. level between them. The prevalence of being “active” stayed within 28% and 30%.

Study	P.A. Level		Chi-square Test <i>p</i> -value
	Some active, Sedentary (%)	Active (%)	
2009	70.7	29.3	0.471
2011	71.6	28.4	

Preliminary Study on the “Sport For All Day”

7.5.16 Close to half of the respondents (46.4%) heard about the arrangement of the “Sport for All Day”. Significant difference was observed among different age groups that 53.7% of young adults heard about the event, 22.9 percent points higher than older adults as shown in the table below.

Heard about the “Sport For All Day” Before	Age (%)			Total (%)	Chi-square Test <i>p</i> -value
	20–39	40–59	60–69		
Yes	53.7	44.1	30.8	46.4	0.000 [@]
No/ Not sure	46.3	55.9	69.2	53.6	

[@] significant difference at $p \leq 0.05$

7.5.17 12.2% of the respondents who were aware of the “Sport for All Day” participated in the event. No significant difference was observed among different age groups as shown in the table below.

Participation in the “Sport For All Day”	Age (%)			Total (%)	Chi-square Test <i>p</i> -value
	20–39	40–59	60–69		
Yes	13.7	10.8	10.5	12.2	0.513
No/ Not sure	86.3	89.2	89.5	87.8	

7.5.18 As much as 60.4% of “Sport for All Day” participants thought that their interests in sports and P.A. could be raised through the participation. No significant difference was found among different age groups as shown in the table below.

Increase Interests in Sports and P.A.	Age (%)			Total (%)	Chi-square Test <i>p</i> -value
	20–39	40–59	60–69		
Yes	57.3	60.8	81.4	60.4	0.403
No	42.7	39.2	18.6	39.6	

7.6 Discussion & Recommendations

7.6.1 It is important to provide not only all-around education on the relationship between physical activity and health to schoolchildren, but also a regular delivery of related messages, including simple concepts of exercise training, health risks associated with inactivity, etc. For example, it is necessary for all ages to learn how to use BMI for overweight and obese classification, in addition to the idea of central obesity. It can be a breakthrough point to alleviate the rapid expansion of obesity. Another example is regular checking of blood pressure. In this Study, results showed that the prevalence of hypertensive response kept on increasing throughout the adulthood. A sharp rise in having hypertension was also observed in women. However, “*Population Health Survey 2003/2004*” found that less than two-thirds of people (62.9%) had their blood pressure checked by health professionals from year 1999 to 2004. In those who had the blood pressure checking, 21.2% performed it once for more than 12 months. Making people more self-conscious about the universal and comparable standards as well as helping them take the initiative to develop an active lifestyle are always the most effective strategies.

- 7.6.2 In the current study, 81.9% of men and 58.1% of women were currently engaged in employment, including full-time and part-time job. They spent 47.75 hours on average per week. 11.0% of them worked even 60 hours or above. 28.3% of adults and elderly had not retired yet. The workplace nowadays is no doubt a second home for many people. It is suggested that employers provide their employees with simple physical activities in the workplace or arrange regular recreation sports activities to them and their family members. It can help raise employees' P.A. level as well as promoting their sense of belonging to the company.
- 7.6.3 Results showed that at least one person was classified as either of overweight or obese for every two men. Even as high as 40.8% of older women experienced the problem of obesity. According to "American Indicator", 71.6% of adults and elderly did not have sufficient P.A. to sustain healthy body. Enormous economic and medical costs will be incurred if no further actions are taken. Related stakeholders can encourage sedentary people to have moderate-or-above intensity P.A. at least three days a week with accumulation of 30 minutes or above per day. In other words, sedentary people are advised to take a first step to meet "Baseline Indicator". For those who have already met the basic requirement, they were suggested to gradually increase their P.A. level and ultimately achieve 150 minutes moderate-or-above intensity P.A. On the other hand, related stakeholders were also recommended to instill the concepts of energy balancing in the public at various levels, including the implementation of food labeling policy, standardized recommendations for daily intake by authority such as universities and professional dieticians, and exercise training principles instructed by qualified fitness instructors.

- 7.6.4 As much as 66.5% and 48.4% of middle-aged men and women respectively encountered overweight or obesity. The situation raised awareness in the society. It is common that the middle-aged are busy at work and find difficulty in sparing time to take part in P.A., which involves with standardized requirements on the number of participants, time, and venue, etc. Related stakeholders may consider organizing some kind of activities, which only require simple equipment and are not restricted by time, say jogging and walking. Other possible activities suitable for middle-aged people include gateball, lawn bowls, hydro-robics, and stretching exercise. The middle-aged can alleviate pressure from work and daily life via making full of the opportunities to engage in more physical activities. On the other hand, it is suggested that business companies design some fitness areas for exercise so as to help employees incorporate a regular exercise habit into workplace.
- 7.6.5 In adults and elderly, “walking”, “running/ jogging”, “ball games”, “swimming”, and “hiking” were the top five popular sports activities. Exercise promotion programs such as walking fun days and family ball games carnivals can be organized periodically at the venues managed by related stakeholders.
- 7.6.6 Evidence showed that the general fitness level of women declined much faster than men across their adulthood in the current study. It is necessary to promote healthy and active lifestyle to women. Related stakeholders may consider organizing more exercise and recreational programs that are appealing to women, such as dancing, hydro-robics, TaiChi, fit ball, Thera-band exercise ...etc.

8 Conclusion

- 8.1** Results showed that different age groups had certain proportions being classified as underweight and overweight/ obese. The situation raised our awareness. According to “Global Strategy on Diet, Physical Activity and Health” released by WHO in 2004, regular P.A. and balanced diet had greater influence on human health. They both helped control the calories intake and retrieve the problem of obesity.

- 8.2** Educators were suggested to take reference from the guidelines prepared by EDB and the Department of Health to improve the inactivity in children. A Physical Education curriculum covering a great variety of physical activity through different learning experiences was needed to provide students with more options. Moreover, it was recommended that related stakeholders offer education to parents who had the responsibility to encourage and support their children to participate in P.A. Community resources could also be further utilized so as to enrich the opportunities of non-school based physical activities for children.
- 8.3** Men generally performed better physical fitness level than women. After reaching adulthood, both genders experienced significant drops in fitness parameters but expansions of health risks (obesity, high BP, and low cardiovascular fitness) as age increased. On the other hand, women were found to have a diminution in fitness at a significantly faster rate than men. Related stakeholders were suggested to strength education and publicity in order to enhance people's cognition in physical fitness as well as their alertness against health risks.
- 8.4** Living habits, such as sleeping time, screen time, and parents' participation frequencies in P.A. were shown to correlate with P.A. level at different degrees of significance. To increase people's fitness level and P.A. level, it was therefore necessary to modify personal lifestyle carefully, say, sufficient sleeping hours (not too many or too few), reducing sedentary activities, increasing involvement in family-based physical activities, maintaining a balanced diet from young, and so on.
- 8.5** Results showed that high correlations only existed between limited numbers of fitness parameters. In other words, every parameter had its unique characteristics and measurement value. It was recommended to include as more test items as possible so as to project a realistic fitness profile.

- 8.6** Hong Kong people were encouraged to gradually develop an active lifestyle starting from “Baseline Indicator” which emphasized the participation in moderate-or-above intensity P.A. at least three days a week with accumulation of 30 minutes or above per day. Research team also pointed out a dose-effect relationship between P.A. levels and health benefits. Everyone should therefore avoid inactivity. Having limited physical activities were stills said to be better than nothing done. Participation in any P.A. would always bring you some health benefits. The long-term goal was to achieve recommendations stated in “American Indicator”. That is, for children, to accumulate at least 60 minutes moderate-or-above intensity P.A. every day in a week; for adolescents, to accumulate at least 60 minutes moderate-or-above intensity P.A. every day in a week in which any three days involved with vigorous intensity P.A.; for adults and elderly, to accumulate at least 150 minutes moderate-or-above intensity P.A. in a week (or 75minutes vigorous intensity P.A. or other combinations of different intensities and durations). People who had not taken part in P.A. for some time, and those with chronic diseases or being at high risks for these problems should consult their doctors before embarking on an exercise training programme.
- 8.7** To conclude, findings indicate no significant rise but a slight drop in the overall P.A. level of Hong Kong people (except for girls aged 7 to 12) in the past two years according to “Baseline Indicator” and “American Indicator”. One of the reasons may be attributed to the occurrence of major sports events. The data collection period of “Sports for All Study 2009” was mainly in year 2008 in which Beijing hosted Olympic Games and Hong Kong co-organized the equestrian competition. At the same time, Hong Kong also actively prepared for the first regional multi-sport event “2009 East Asian Games”. The government devoted resources to popularize the event with the support of the mass media. The passionate atmosphere for physical activities was developed as a result. People may easily get in touch with the exercise and sports as well as yielding interests in participation, raising the P.A. level. Since the Games were all over, the atmosphere cooled down and the P.A. level may decrease eventually.

8.8 Since 1995, the Government of the People's Republic of China announced the National policy “全民健身計劃綱要” and has determined “Fitness for All” as their national health objective. Since then significant education and researches are actively implemented throughout the country. The United States has established a government funded and independent agency called the “National Coalition for Promoting Physical Activity (NCPA)” as a result of the release of the Surgeon General Report on Physical Activity and Health (1996). Increasing the P.A. level became the top priority of their national health goal, followed by obesity reduction and smoking cessation. Professional bodies like the American College of Sports Medicine, American Heart Association, and President Challenge for Physical Fitness Council jointed the NCPA for implementing nationwide P.A. promotion to emphasize the importance of regular P.A. To learn from these experiences, we may set citywide health objectives in Hong Kong and put increasing the P.A. level as one of our top priority. The Study is the first scientific-based survey which aimed to investigate Hong Kong people's fitness level and P.A. participation pattern using random sampling. It provided an enormous database to help define the directions and strategies. Once objectives are established, evaluation can be easily made in future to see whether our existing resources are effective to achieve the health objectives.

- END -